

**AMENDMENTS TO THE CLAIMS**

1-2. (Cancelled)

3. (Currently amended) A method for inhibition of keloid and/or hypertrophic scar formation in a course of therapy of wound or dermal injury and for ~~inhibition-treatment~~ of pain or pruritus on ~~the a wound~~ lesion of keloid and/or hypertrophic scar which comprises administering a medicament containing acetylsalicylic acid ~~or its pharmacologically acceptable salt~~ in an effective amount to said wound lesion of a patient.

4. (Cancelled)

5. (Currently amended) A method for inhibition of keloid and/or hypertrophic scar formation in a course of therapy of wound or dermal injury which comprises administering a medicament containing acetylsalicylic acid ~~or its pharmacologically acceptable salt~~ in an effective amount to a wound lesion of a patient.

6. (New) A method for inhibition of keloid and/or hypertrophic scar formation in a course of therapy of wound or dermal injury which comprises administering a medicament containing acetylsalicylic acid in an effective amount of 0.005 to 1000  $\mu\text{g/g}$  per tissue weight to a wound lesion of a patient.

7. (New) A method for inhibition of keloid and/or hypertrophic scar formation in a course of therapy of wound or dermal injury which comprises administering a medicament containing acetylsalicylic acid in an effect amount of 0.01 to 800  $\mu\text{g/g}$  per tissue weight to a wound lesion of a patient.